

A CONSTRUCTION OF HORIKAWA SURFACE VIA \mathbb{Q} -GORENSTEIN SMOOTHINGS

YONGNAM LEE AND JONGIL PARK

ABSTRACT. In this article we prove that Fintushel-Stern's construction of Horikawa surface, which is obtained from an elliptic surface via a rational blow-down surgery in smooth category, can be performed in complex category. The main technique involved is \mathbb{Q} -Gorenstein smoothings.

1. INTRODUCTION

As an application of a rational blow-down surgery on 4-manifolds, R. Fintushel and R. Stern showed that Horikawa surface $H(n)$ can be obtained from an elliptic surface $E(n)$ via a rational blow-down surgery in smooth category [3]. Note that Horikawa surface $H(n)$ is defined as a double cover of a Hirzebruch surface \mathbb{F}_{n-3} branched over $|6C_0 + (4n - 8)f|$, where C_0 is a negative section and f is a fiber of \mathbb{F}_{n-3} .

In this article we show that a rational blow-down surgery to obtain Horikawa surface can be performed in fact in complex category. That is, we reinterpret algebraically Fintushel-Stern's topological construction [3] of Horikawa surface $H(n)$ to give a complex structure on it. The main technique we use in this paper is \mathbb{Q} -Gorenstein smoothings. Note that \mathbb{Q} -Gorenstein smoothing theory developed in deformation theory in last thirty years is a very powerful tool to construct a non-singular surface of general type. The basic scheme is the following: Suppose that a projective surface contains several disjoint chains of curves representing the resolution graphs of special quotient singularities. Then, by contracting these chains of curves, we get a singular surface X with special quotient singularities. And then we investigate the existence of a \mathbb{Q} -Gorenstein smoothing of X . It is known that the cohomology $H^2(T_X^0)$ contains the obstruction space of a \mathbb{Q} -Gorenstein smoothing of X . That is, if $H^2(T_X^0) = 0$, then there is a \mathbb{Q} -Gorenstein smoothing of X . For example, we recently constructed a simply connected minimal surface of general type with $p_g = 0$ and $K^2 = 2$ by proving the cohomology $H^2(T_X^0) = 0$ [9]. But, in general, the cohomology $H^2(T_X^0)$ is not zero and it is a very difficult problem to determine whether there exists a \mathbb{Q} -Gorenstein smoothing of X . In this article we also give a family of examples which admit \mathbb{Q} -Gorenstein smoothings even though the cohomology $H^2(T_X^0)$ does not vanish. Our main technique is a \mathbb{Q} -Gorenstein smoothing theory with a cyclic group action. It is briefly reviewed and developed in Section 2.

The sketch of our construction whose details are given in Section 3 is as follows: We first construct a simply connected relatively minimal elliptic surface $E(n)$ ($n \geq 5$) with

Date: June 1, 2008.

2000 Mathematics Subject Classification. Primary 14J29; Secondary 14J10, 14J17.

Key words and phrases. Horikawa surface, \mathbb{Q} -Gorenstein smoothing, rational blow-down.

a special fiber, which contains two linear chains of configurations of \mathbb{P}^1 's

$$\begin{array}{ccccccc} \overset{-n}{\circ} & - & \overset{-2}{\circ} & - & \overset{-2}{\circ} & - & \cdots - & \overset{-2}{\circ} \\ U_{n-3} & & U_{n-4} & & U_{n-5} & & & U_1 \end{array}.$$

We construct this kind of an elliptic surface $E(n)$ explicitly by using a double cover of a blowing-up of Hirzebruch surface \mathbb{F}_n branched over a special curve. The double cover of \mathbb{F}_n has two rational double points A_1 and A_{2n-9} . Then its minimal resolution is an elliptic surface $E(n)$ which has an I_{2n-6} as a special fiber. Now we contract these two linear chains of configurations of \mathbb{P}^1 's to produce a normal projective surface X_n with two special quotient singularities, both singularities are of type $\frac{1}{(n-2)^2}(1, n-3)$. Finally, we apply \mathbb{Q} -Gorenstein smoothing theory with a cyclic group action developed in Section 2 for X_n in order to get our main result which is following.

Theorem 1.1. *The projective surface X_n obtained by contracting two disjoint configurations C_{n-2} from an elliptic surface $E(n)$ admits a \mathbb{Q} -Gorenstein smoothing of two quotient singularities all together, and a general fiber of the \mathbb{Q} -Gorenstein smoothing is Horikawa surface $H(n)$.*

Acknowledgements. The authors would like to thank Ronald Fintushel and Ronald Stern for helpful comments on Horikawa surfaces during the Conference of Algebraic Surfaces and 4-manifolds held at KIAS. The authors also wish to thank Roberto Pignatelli for valuable discussions to prove Proposition 3.1. Yongnam Lee was supported by Korea Research Foundation Grant funded by the Korean Government (KRF-2005-070-C00005). He would like to thank Meng Chen for his generous hospitality during his visit Fudan University, where some of the paper was completed. Jongil Park was supported by SBS Foundation Grant in 2007 and he also holds a joint appointment in the Research Institute of Mathematics, Seoul National University.

2. \mathbb{Q} -GORENSTEIN SMOOTHING

In this section we briefly review a theory of \mathbb{Q} -Gorenstein smoothing for projective surfaces with special quotient singularities, which is a key technical ingredient in our main construction.

Definition. Let X be a normal projective surface with quotient singularities. Let $\mathcal{X} \rightarrow \Delta$ (or \mathcal{X}/Δ) be a flat family of projective surfaces over a small disk Δ . The one-parameter family of surfaces $\mathcal{X} \rightarrow \Delta$ is called a *\mathbb{Q} -Gorenstein smoothing* of X if it satisfies the following three conditions;

- (i) the general fiber X_t is a smooth projective surface,
- (ii) the central fiber X_0 is X ,
- (iii) the canonical divisor $K_{\mathcal{X}/\Delta}$ is \mathbb{Q} -Cartier.

A \mathbb{Q} -Gorenstein smoothing for a germ of a quotient singularity $(X_0, 0)$ is defined similarly. A quotient singularity which admits a \mathbb{Q} -Gorenstein smoothing is called a *singularity of class T* .

Proposition 2.1 ([8, 12, 16]). *Let $(X_0, 0)$ be a germ of two dimensional quotient singularity. If $(X_0, 0)$ admits a \mathbb{Q} -Gorenstein smoothing over the disk, then $(X_0, 0)$ is either a rational double point or a cyclic quotient singularity of type $\frac{1}{dn^2}(1, dna-1)$ for some integers a, n, d with a and n relatively prime.*

Proposition 2.2 ([8, 12, 17]). (1) The singularities $\overset{-4}{\circ}$ and $\overset{-3}{\circ} - \overset{-2}{\circ} - \overset{-2}{\circ} - \dots - \overset{-2}{\circ} - \overset{-3}{\circ}$ are of class T .

(2) If the singularity $\overset{-b_1}{\circ} - \dots - \overset{-b_r}{\circ}$ is of class T , then so are

$$\overset{-2}{\circ} - \overset{-b_1}{\circ} - \dots - \overset{-b_{r-1}}{\circ} - \overset{-b_r-1}{\circ} \quad \text{and} \quad \overset{-b_1-1}{\circ} - \overset{-b_2}{\circ} - \dots - \overset{-b_r}{\circ} - \overset{-2}{\circ}.$$

(3) Every singularity of class T that is not a rational double point can be obtained by starting with one of the singularities described in (1) and iterating the steps described in (2).

Let X be a normal projective surface with singularities of class T . Due to the result of Kollár and Shepherd-Barron [8], there is a \mathbb{Q} -Gorenstein smoothing locally for each singularity of class T on X (see Proposition 2.5). The natural question arises whether this local \mathbb{Q} -Gorenstein smoothing can be extended over the global surface X or not. Roughly geometric interpretation is the following: Let $\cup_{\alpha} V_{\alpha}$ be an open covering of X such that each V_{α} has at most one singularity of class T . By the existence of a local \mathbb{Q} -Gorenstein smoothing, there is a \mathbb{Q} -Gorenstein smoothing $\mathcal{V}_{\alpha}/\Delta$. The question is if these families glue to a global one. The answer can be obtained by figuring out the obstruction map of the sheaves of deformation $T_X^i = \text{Ext}_X^i(\Omega_X, \mathcal{O}_X)$ for $i = 0, 1, 2$. For example, if X is a smooth surface, then T_X^0 is the usual holomorphic tangent sheaf T_X and $T_X^1 = T_X^2 = 0$. By applying the standard result of deformations [10, 14] to a normal projective surface with quotient singularities, we get the following

Proposition 2.3 ([16], §4). Let X be a normal projective surface with quotient singularities. Then

- (1) The first order deformation space of X is represented by the global Ext 1-group $\mathbb{T}_X^1 = \text{Ext}_X^1(\Omega_X, \mathcal{O}_X)$.
- (2) The obstruction lies in the global Ext 2-group $\mathbb{T}_X^2 = \text{Ext}_X^2(\Omega_X, \mathcal{O}_X)$.

Furthermore, by applying the general result of local-global spectral sequence of ext sheaves ([14], §3) to deformation theory of surfaces with quotient singularities so that $E_2^{p,q} = H^p(T_X^q) \Rightarrow \mathbb{T}_X^{p+q}$, and by $H^j(T_X^i) = 0$ for $i, j \geq 1$, we also get

Proposition 2.4 ([12, 16]). Let X be a normal projective surface with quotient singularities. Then

- (1) We have the exact sequence

$$0 \rightarrow H^1(T_X^0) \rightarrow \mathbb{T}_X^1 \rightarrow \ker[H^0(T_X^1) \rightarrow H^2(T_X^0)] \rightarrow 0$$

where $H^1(T_X^0)$ represents the first order deformations of X for which the singularities remain locally a product.

- (2) If $H^2(T_X^0) = 0$, every local deformation of the singularities may be globalized.

The vanishing $H^2(T_X^0) = 0$ can be obtained via the vanishing of $H^2(T_V(-\log E))$, where V is the minimal resolution of X and E is the reduced exceptional divisors. Note that every singularity of class T has a local \mathbb{Q} -Gorenstein smoothing by Proposition 2.5 below.

Let X be a normal projective surface with singularities of class T . Our concern is to understand \mathbb{Q} -Gorenstein smoothings in \mathbb{T}_X^1 , not the whole first order deformations. These special deformations can be constructed via local index one cover. Let $U \subset X$ be an analytic neighborhood with an index one cover U' . For the case of the field \mathbb{C} ,

this index one cover is unique up to isomorphism. The first order deformations which associate \mathbb{Q} -Gorenstein smoothings can be realized as the invariant part of $T_{\mathcal{U}'}^1$. The sheaves \tilde{T}_X^1 are defined by the index one covering stack and by the étale sites [5]. The first order deformation of a \mathbb{Q} -Gorenstein smoothing of singularities of class T is expressed by the cohomology $H^0(\tilde{T}_X^1)$ [5, 7, 8]. By the help of the birational geometry in threefolds and their applications to deformations of surface singularities, the following proposition is obtained. Note that the cohomology $H^0(\tilde{T}_X^1)$ is given explicitly as follows.

Proposition 2.5 ([8, 12]). (1) *Let $a, d, n > 0$ be integers with a, n relatively prime and consider a map $\pi : \mathcal{Y}/\mu_n \rightarrow \mathbb{C}^d$, where $\mathcal{Y} \subset \mathbb{C}^3 \times \mathbb{C}^d$ is the hypersurface of equation $uv - y^{dn} = \sum_{k=0}^{d-1} t_k y^{kn}$; t_0, \dots, t_{d-1} are linear coordinates over \mathbb{C}^d , μ_n acts on \mathcal{Y} by*

$$\mu_n \ni \xi : (u, v, y, t_0, \dots, t_{d-1}) \rightarrow (\xi u, \xi^{-1} v, \xi^a y, t_0, \dots, t_{d-1})$$

and π is the factorization to the quotient of the projection $\mathcal{Y} \rightarrow \mathbb{C}^d$. Then π is a \mathbb{Q} -Gorenstein smoothing of the cyclic singularity of a germ $(X_0, 0)$ of type $\frac{1}{dn^2}(1, dna-1)$. Moreover every \mathbb{Q} -Gorenstein smoothing of $(X_0, 0)$ is isomorphic to the pull-back of π for some germ of holomorphic map $(\mathbb{C}, 0) \rightarrow (\mathbb{C}^d, 0)$.

(2) *Let X be a normal projective surface with singularities of class T . Then*

$$H^0(\tilde{T}_X^1) = \sum_{p \in \text{singular points of } X} \mathbb{C}_p^{\oplus d_p}$$

where a singular point p is of type $\frac{1}{d_p n^2}(1, d_p a n - 1)$ with $(a, n) = 1$.

Theorem 2.1 ([9]). *Let X be a normal projective surface with singularities of class T . Let $\pi : V \rightarrow X$ be the minimal resolution and let E be the reduced exceptional divisors. Suppose that $H^2(T_V(-\log E)) = 0$. Then $H^2(T_X^0) = 0$ and there is a \mathbb{Q} -Gorenstein smoothing of X .*

As we see in Theorem 2.1 above, if $H^2(T_X^0) = 0$, then there is a \mathbb{Q} -Gorenstein smoothing of X . For example, we constructed a simply connected minimal surface of general type with $p_g = 0$ and $K^2 = 2$ by proving the cohomology $H^2(T_X^0) = 0$ [9]. But, in general, the cohomology $H^2(T_X^0)$ is not zero and it is a very difficult problem to determine whether there exists a \mathbb{Q} -Gorenstein smoothing of X . Hence, in the case that $H^2(T_X^0) \neq 0$, we have to develop another technique in order to investigate the existence of \mathbb{Q} -Gorenstein smoothings. Even though we do not know whether such a technique exists in general, if X is a normal projective surface with singularities of class T which admits a cyclic group with some nice properties, then we are able to show that it admits a \mathbb{Q} -Gorenstein smoothing. Explicitly, we get the following theorem.

Theorem 2.2. *Let X be a normal projective surface with singularities of class T . Assume that a cyclic group G acts on X such that*

- (1) $Y = X/G$ is a normal projective surface with singularities of T ,
- (2) $p_g(Y) = q(Y) = 0$,
- (3) Y has a \mathbb{Q} -Gorenstein smoothing,
- (4) the map $\sigma : X \rightarrow Y$ induced by a cyclic covering is flat, and the branch locus D (resp. the ramification locus) of the map $\sigma : X \rightarrow Y$ is an irreducible nonsingular curve lying outside the singular locus of Y (resp. of X), and
- (5) $H^1(Y, \mathcal{O}_Y(D)) = 0$.

Then there exists a \mathbb{Q} -Gorenstein smoothing of X that is compatible with a \mathbb{Q} -Gorenstein smoothing of Y . And the cyclic covering extends to the \mathbb{Q} -Gorenstein smoothing.

Proof. Let $\mathcal{Y} \rightarrow \Delta$ be a \mathbb{Q} -Gorenstein smoothing of Y , and let Y_t be a general fiber of the \mathbb{Q} -Gorenstein smoothing. By the semi-continuity, we have $p_g(Y_t) = q(Y_t) = 0$. The base change theorem and Leray spectral sequence imply that $H^1(\mathcal{Y}, \mathcal{O}_{\mathcal{Y}}) = H^2(\mathcal{Y}, \mathcal{O}_{\mathcal{Y}}) = 0$. It gives an isomorphism $r_0 : \text{Pic}(\mathcal{Y}) \rightarrow \text{Pic}(Y)$ and an injective map $r_t : \text{Pic}(\mathcal{Y}) \rightarrow \text{Pic}(Y_t)$ (Lemma 2 in [12]). The vanishing $H^1(Y, \mathcal{O}_Y(D)) = 0$ ensures that the deformation of Y can be lifted to the deformation of the pair (Y, D) , i.e. the branch divisor D is extended to D_t in Y_t . Since the divisor D is nonsingular, D_t is also nonsingular. And the flatness of the map ensures that the divisor L which is the data of the cyclic cover, i.e. $L^{\otimes |G|} \cong D$, is extended to L_t with $L_t^{\otimes |G|} \cong D_t$. Hence, the cyclic covering extends to the \mathbb{Q} -Gorenstein smoothing of Y . \square

3. A CONSTRUCTION OF HORIKAWA SURFACE

Let $E(n)$ be a simply connected relatively minimal elliptic surface with a section and with $c_2 = 12n$. Then there is only one up to diffeomorphism such an elliptic surface and the canonical class is given by $K_{E(n)} = (n-2)C$, where C is a general fiber of an elliptic fibration. Hence each section is a nonsingular rational curve whose self-intersection number is $-n$. Assume that $n \geq 4$ and let C_{n-2} denote a simply connected smooth 4-manifold obtained by plumbing the $(n-3)$ disk bundles over the 2-sphere according to the linear diagram

$$\begin{array}{ccccccc} \overset{-n}{\circ} & - & \overset{-2}{\circ} & - & \overset{-2}{\circ} & - & \cdots - & \overset{-2}{\circ} \\ U_{n-3} & & U_{n-4} & & U_{n-5} & & & U_1 \end{array}$$

Assume that an elliptic surface $E(n)$ has two configurations C_{n-2} such that all embedded 2-spheres U_i are holomorphic curves (We show the existence of such an $E(n)$ later). Let Y'_n be a normal projective surface obtained by contracting one configuration C_{n-2} from $E(n)$. Then Y'_n does not admit a \mathbb{Q} -Gorenstein smoothing because it violates Noether inequality (Corollary 7.5 in [3]). In fact, it does not satisfy the vanishing condition in the hypothesis of Theorem 2.1, that is, we have $H^2(E(n), T_{E(n)}) \neq 0$: Let $h : E(n) \rightarrow \mathbb{P}^1$ be an elliptic fibration. Assume that C is a general fiber of the map h . We have an injective map $0 \rightarrow h^*\Omega_{\mathbb{P}^1} \rightarrow \Omega_{E(n)}$ and the map induces an injection $H^0(\mathbb{P}^1, \Omega_{\mathbb{P}^1}(n-2)) \hookrightarrow H^0(E(n), \Omega_{E(n)}((n-2)C))$ by tensoring $(n-2)C$ on $0 \rightarrow h^*\Omega_{\mathbb{P}^1} \rightarrow \Omega_{E(n)}$. Since $K_{E(n)} = (n-2)C$, the cohomology $H^0(E(n), \Omega_{E(n)}(K_{E(n)}))$ is not zero. Hence the Serre duality implies that $H^2(E(n), T_{E(n)})$ is not zero.

Next, let X_n be a normal projective surface obtained by contracting two disjoint configurations C_{n-2} from $E(n)$, and we want to investigate the existence of a \mathbb{Q} -Gorenstein smoothing of X_n . As a warming-up, we first investigate $n = 4$ case.

Example. R. Gompf constructed a family of symplectic 4-manifolds by taking a fiber sum of other symplectic 4-manifolds [4]. To recall Gompf's example briefly, we start with a simply connected relatively minimal elliptic surface $E(4)$ with a section and with $c_2 = 48$. It is known that $E(4)$ admits nine rational (-4) -curves as disjoint sections. Rationally blowing-down n (-4) -curves of $E(4)$ is the same as the normal connected sum of $E(4)$ with n copies of \mathbb{P}^2 by identifying a conic in each \mathbb{P}^2 with one (-4) -curve in $E(4)$. Let us denote this 4-manifold by $W_{4,n}$. Then the manifold $W_{4,1}$ does not admit

any complex structure because it violates the Noether inequality. But we will show that $W_{4,2}$ admits a complex structure using a \mathbb{Q} -Gorenstein smoothing theory. For this, let us first denote the singular projective surface obtained by contracting n (-4) -sections from $E(4)$ by $W'_{4,n}$. And then we claim that $W'_{4,2}$ has a \mathbb{Q} -Gorenstein smoothing. The reason is following: Consider $E(4)$ as a double cover of Hirzebruch surface \mathbb{F}_4 branched over an irreducible nonsingular curve D in the linear system $|4(C_0 + 4f)|$, where C_0 is a negative section and f is a fiber of \mathbb{F}_4 . Then $H^1(\mathbb{F}_4, \mathcal{O}_{\mathbb{F}_4}(D)) = 0$: Since $p_g(\mathbb{F}_4) = q(\mathbb{F}_4) = 0$,

$$H^1(\mathbb{F}_4, \mathcal{O}_{\mathbb{F}_4}(D)) \simeq H^1(D, \mathcal{O}_D(D)) \simeq H^0(D, \mathcal{O}_D(K_D - D))^\vee.$$

And $\deg K_D - D^2 = 4(C_0 + 4f)(2C_0 + 10f) - 16(C_0 + 4f)^2 = -24 < 0$ implies that $H^0(D, \mathcal{O}_D(K_D - D)) = 0$. Since D does not intersect C_0 , $W'_{4,2}$ is a double cover of a cone $\hat{\mathbb{F}}_4$ which is a contraction of C_0 from \mathbb{F}_4 . This implies that the map σ induced by a double cover is flat and $H^1(Y, \mathcal{O}_Y(D)) = 0$. Note that $\hat{\mathbb{F}}_4$ has a \mathbb{Q} -Gorenstein smoothing whose general fiber is \mathbb{P}^2 . It is obtained by a pencil of hyperplane section of the cone of the Veronese surface imbedded in \mathbb{P}^5 . Hence $W'_{4,2}$ has a \mathbb{Q} -Gorenstein smoothing by Theorem 2.2. Finally, since the rational blow-down manifold $W_{4,2}$ is diffeomorphic to the general fiber of the \mathbb{Q} -Gorenstein smoothing of $W'_{4,2}$, $W_{4,2}$ admits a complex structure. Furthermore, using a triple cover of \mathbb{F}_4 branched over D in the linear system $|3(C_0 + 4f)|$, we can also prove that $W'_{4,3}$ has a \mathbb{Q} -Gorenstein smoothing by the similar proof as above. And, by extending Theorem 2.2 to a finite abelian group, it is possible to show that some other manifolds $W'_{4,n}$ has a \mathbb{Q} -Gorenstein smoothing, too. We leave it for a future research.

Now we investigate the general case. Assume that $n \geq 5$ and let \mathbb{F}_n be a Hirzebruch surface. Let C_0 be a negative section with $C_0^2 = -n$ and f be a fiber of \mathbb{F}_n . Consider the linear system $|4(C_0 + nf)|$. The surface \mathbb{F}_n can be obtained from the cone over a rational normal curve of degree n by blowing up the vertex. And a curve in the linear system $|4(C_0 + nf)|$ is the strict transform of the hyperplane section of the cone. By Bertini's theorem, there is an irreducible nonsingular curve in the linear system $|4(C_0 + nf)|$. The double cover of \mathbb{F}_n branched over an irreducible nonsingular member in $|4(C_0 + nf)|$ is an elliptic surface $E(n)$: Let $\sigma : \hat{X}_n \rightarrow \mathbb{F}_n$ be a double covering branched over an irreducible nonsingular member in the linear system $|4(C_0 + nf)|$. Then, by the invariants of a double covering ([2], Chapter V), we have $p_g(\hat{X}_n) = p_g(\mathbb{F}_n) + h^0(\mathbb{F}_n, K_{\mathbb{F}_n} + L) = h^0(\mathbb{F}_n, (n - 2)f) = n - 1$ and $\chi(\mathcal{O}_{\hat{X}_n}) = 2\chi(\mathcal{O}_{\mathbb{F}_n}) + \frac{1}{2}(L \cdot K_{\mathbb{F}_n}) + \frac{1}{2}(L \cdot L) = n$, where $L = 2(C_0 + nf)$. Therefore we have $q(\hat{X}_n) = 0$ and $K_{\hat{X}_n}^2 = 2(\sigma^*(K_{\mathbb{F}_n} + L))^2 = 2((n - 2)f)^2 = 0$.

In this article we want to choose a special irreducible (singular) curve D in the linear system $|4(C_0 + nf)|$, which has a special intersection with one special fiber f : Note that $D \cdot f = 4$. We want D to intersect with f at two distinct points p and q that are not in C_0 . Let $x = 0$ be the local equation of f and x, y be a coordinate at p (resp. at q). We require that the local equation of D at p (resp. at q) is $(y - x)(y + x) = 0$ (resp. $(y - x^{n-4})(y + x^{n-4}) = 0$). These are $3(n - 4) + 3$ -conditions: $1, x, x^2, \dots, x^{2n-9}, y, yx, \dots, yx^{n-5}$ terms should vanish to have the local analytic equation $(y - x^{n-4})(y + x^{n-4}) = 0$. By next lemmas and proposition, we have such a curve D satisfying the conditions above.

Lemma 3.1. *We have $h^0(\mathbb{F}_n, \mathcal{O}_{\mathbb{F}_n}(D)) = 10n + 5$, where D is a member in the linear system $|4(C_0 + nf)|$.*

Proof. Let $C = C_0 + nf$. Then, by the following two exact sequences

$$0 \rightarrow \mathcal{O}_{\mathbb{F}_n} \rightarrow \mathcal{O}_{\mathbb{F}_n}(f) \rightarrow \mathcal{O}_{\mathbb{P}^1} \rightarrow 0$$

$$0 \rightarrow \mathcal{O}_{\mathbb{F}_n}(kf) \rightarrow \mathcal{O}_{\mathbb{F}_n}((k+1)f) \rightarrow \mathcal{O}_{\mathbb{P}^1} \rightarrow 0,$$

we have $h^1(\mathbb{F}_n, \mathcal{O}_{\mathbb{F}_n}(kf)) = 0$ for all nonnegative integers k . And from the exact sequence

$$0 \rightarrow \mathcal{O}_{\mathbb{F}_n}(nf) \rightarrow \mathcal{O}_{\mathbb{F}_n}(C) \rightarrow \mathcal{O}_{C_0}(C) \rightarrow 0,$$

we also have $h^1(\mathbb{F}_n, \mathcal{O}_{\mathbb{F}_n}(C)) = 0$ and $h^0(\mathbb{F}_n, \mathcal{O}_{\mathbb{F}_n}(C)) = n + 2$. Hence, by considering the exact sequences similarly

$$0 \rightarrow \mathcal{O}_{\mathbb{F}_n}(kC) \rightarrow \mathcal{O}_{\mathbb{F}_n}((k+1)C) \rightarrow \mathcal{O}_C((k+1)C) \rightarrow 0,$$

we finally get $h^0(\mathbb{F}_n, \mathcal{O}_{\mathbb{F}_n}(D)) = 10n + 5$. \square

First, we assume that D is nonsingular at every point except the two points p and q . Let $\sigma : \tilde{X}_n \rightarrow \mathbb{F}_n$ be a double covering branched over the curve D chosen above. Then \tilde{X}_n is a singular elliptic surface with $p_g = n - 1$ and $\chi = n$ which has two rational double points by the local equations of D at p and q - one is A_1 ($z^2 = y^2 - x^2$) and the other one is A_{2n-9} ($z^2 = y^2 - x^{2n-8}$). Therefore its minimal resolution is also an elliptic surface $E(n)$. First we blow up at p and q in \mathbb{F}_n . Then we have an exceptional curve coming from a blowing up at p which intersects with the proper transform of D transversally at two points, and we also have an exceptional curve coming from a blowing up at q which intersects with the proper transform of D at one point, say q_1 . Let $x = 0$ be the local equation of the (-1) -exceptional curve at q_1 . Then the local equation of the proper transform of D at q_1 is $(y - x^{n-5})(y + x^{n-5}) = 0$. We blow up again at q_1 . By the continuation of blowing up at infinitely near points of q , we have the following configuration of smooth rational curves

$$\begin{array}{ccccccc} \overset{-n}{\circ} & - & \overset{-2}{\circ} & - & \overset{-2}{\circ} & - \cdots - & \overset{-2}{\circ} \\ U_{n-3} & & U_{n-4} & & U_{n-5} & & U_1 \\ & & | & & & & | \\ & & \circ & & & & \circ \\ & & E_{1,-1} & & & & E_{2,-1} \end{array}$$

where the proper transform of D intersects with E_i , $i = 1, 2$ at two points transversally. We denote this surface by Z_n , which is obtained by $(n - 3)$ times blowing-ups of \mathbb{F}_n .

Let $\pi : Z_n \rightarrow \mathbb{F}_n$ be a map and $\Delta = \pi^*(4C_0 + 4nf) - 2E_1 - 2U_{n-5} - 4U_{n-6} - 6U_{n-7} - \cdots - 2(n-5)U_1 - 2(n-4)E_2$. For a simple computation, we write it as

$$\Delta = \pi^*(4C_0 + 4nf) - 2F - 2 \sum_{i=1}^{n-4} F_i,$$

where $F = E_1$, $F_1 = U_{n-5} + U_{n-6} + \cdots + E_2$, $F_2 = U_{n-6} + \cdots + E_2, \dots, F_{n-4} = E_2$. Note that F_i is not necessarily irreducible and $F^2 = F_i^2 = K_{Z_n} \cdot F = K_{Z_n} \cdot F_i = -1$ for all $i = 1, \dots, n-4$. Let $f_0 = U_{n-4}$, which is a proper transform of the fiber, and let $L = \Delta - (\pi^*C_0 + f_0) - K_{Z_n}$.

In Proposition 3.1 below, we prove that the linear system $|\Delta|$ is base point free. Then, by Bertini's theorem, we conclude that D is nonsingular except the two points p and q .

Lemma 3.2. $L^2 \geq 5$ and L is nef.

Proof. Since $\pi^*f = f_0 + F + \sum_{i=1}^{n-4} F_i$, we have $L = \pi^*(5C_0 + (5n+2)f) - f_0 - 3F - 3\sum_{i=1}^{n-4} F_i = \pi^*(5C_0 + (5n-1)f) + 2f_0$. Furthermore, since the two points p and q are not in C_0 , we also have $C_0 \cdot f = C_0 \cdot f_0 = 1$ and $f_0^2 = -2$. Therefore

$$L^2 = -25n + (5n-1)10 + 20 - 8 = 25n + 2 \geq 5.$$

Let G be an irreducible curve which is neither f_0 nor C_0 . Note that $L \cdot f_0 = 5 - 4 = 1$ and $L \cdot C_0 = -5n + 5n - 1 + 2 = 1$. Write $\pi_*G = aC_0 + bf$. Then we have

$$G \cdot L \geq \pi_*G \cdot (5C_0 + (5n-1)f) = (aC_0 + bf) \cdot (5C_0 + (5n-1)f) = -a + 5b.$$

We note that the linear system $|aC_0 + bf|$ contains an irreducible curve in \mathbb{F}_n if and only if $a = 0, b = 1$; or $a = 1, b = 0$; or $a > 0, b > an$; or $a > 0, b = an$ with $n > 0$ (refer to Corollary 2.18, Chapter V in [6]). Therefore it is impossible that $G \cdot L < 0$. It implies that L is nef. \square

Lemma 3.3. *The linear system $|\Delta - (\pi^*C_0 + f_0)|$ on Z_n is base point free.*

Proof. By Lemma 3.2 above, L is nef and $L^2 \geq 5$. Hence, applying to Reider's theorem [15], if the adjoint linear series $|\Delta - (\pi^*C_0 + f_0)| = |K_{Z_n} + L|$ has a base point at x then there is an effective divisor G in Z_n passing through x such that either $G \cdot L = 0$ and $G^2 = -1$; or $G \cdot L = 1$ and $G^2 = 0$.

Assume that $G \cdot L = 0$ and $G^2 = -1$. Write $G = G_1 + \cdots + G_k$, where G_k is an irreducible curve. Since L is nef, $G \cdot L = 0$ implies that $G_i \cdot L = 0$ for all $i = 1, \dots, k$. Then we get a contradiction by a similar argument to show that $-a + 5b \leq 0$ in the proof Lemma 3.2 above.

Assume that $G \cdot L = 1$ and $G^2 = 0$. By the same argument in the case $G \cdot L = 0$, $G = G_1$. Then we get a contradiction by a similar argument to show that $-a + 5b \leq 1$ unless $G = C_0$ or f_0 . Furthermore, since $C_0^2 = -n$ and $f_0^2 = -2$, it also contradicts. \square

Proposition 3.1. *The linear system $|\Delta|$ on Z_n is base point free.*

Proof. Note that $\Delta \cdot \pi^*C_0 = \Delta \cdot f_0 = 0$. Therefore we have $\mathcal{O}_{\pi^*C_0+f_0}(\Delta) = \mathcal{O}_{\pi^*C_0+f_0}$. Furthermore, by Lemma 3.2 above and by the vanishing theorem, we also have

$$H^1(Z_n, \Delta - (\pi^*C_0 + f_0)) = H^1(Z_n, K_{Z_n} + L) = 0.$$

Hence, using Lemma 3.3 above and the short exact sequence

$$0 \rightarrow \mathcal{O}_{Z_n}(\Delta - (\pi^*C_0 + f_0)) \rightarrow \mathcal{O}_{Z_n}(\Delta) \rightarrow \mathcal{O}_{\pi^*C_0+f_0} \rightarrow 0,$$

we conclude that the linear system $|\Delta|$ is base point free. \square

Next, by Artin's criterion of contraction [1], we can contract a configuration C_{n-2} , which is a linear chain of \mathbb{P}^1 's

$$\begin{array}{ccccccc} \overset{-n}{\circ} & & \overset{-2}{\circ} & & \overset{-2}{\circ} & & \overset{-2}{\circ} \\ U_{n-3} & - & U_{n-4} & - & U_{n-5} & - \cdots - & U_1 \end{array}$$

so that it produces a singular normal projective surface. We denote this surface by Y_n . We note that Δ is the proper transform of D in Z_n and that Y_n has a cyclic quotient singularity of type $\frac{1}{(n-2)^2}(1, n-3)$, which is a singularity of class T .

Lemma 3.4. $H^1(Z_n, \mathcal{O}_{Z_n}(\Delta)) = 0$.

Proof. Since $p_g(Z_n) = q(Z_n) = 0$, we have

$$H^1(Z_n, \mathcal{O}_{Z_n}(\Delta)) \simeq H^1(\Delta, \mathcal{O}_\Delta(\Delta)) \simeq H^0(\Delta, \mathcal{O}_\Delta(K_\Delta - \Delta))^\vee.$$

We also have $\Delta^2 = D^2 - 4(n-3) = 12n + 12$ and $\deg K_\Delta = \deg K_D - 2(n-3) = 12n - 8 - 2(n-3) = 10n - 2$. Therefore it satisfies

$$\deg K_\Delta - \Delta^2 = 10n - 2 - 12n - 12 < 0,$$

and it implies that $H^0(\Delta, \mathcal{O}_\Delta(K_\Delta - \Delta)) = 0$. \square

Proposition 3.2. *The singular surface Y_n admits a \mathbb{Q} -Gorenstein smoothing.*

Proof. It is enough to show that $-K_{Y_n}$ is effective (Theorem 21 in [12]). Let $\pi : Z_n \rightarrow \mathbb{F}_n$ be a composition of blowing-ups, and $\psi : Z_n \rightarrow Y_n$ be a contraction. Then we have

$$K_{Z_n} = \pi^* K_{\mathbb{F}_n} + E_1 + U_{n-5} + 2U_{n-6} + \cdots + (n-5)U_1 + (n-4)E_2.$$

Since $K_{\mathbb{F}_n} = -2C_0 - (n+2)f$, $-K_{Z_n}$ is effective. Furthermore, since $h^0(-K_{Y_n}) = h^0(\psi_*(-K_{Z_n})) = h^0(-K_{Z_n})$ (§3.9.2 in [16]), $-K_{Y_n}$ is also effective. \square

Now we are in a position to prove our main theorem mentioned in the Introduction. First remind that Horikawa surface $H(n)$ is a double cover of \mathbb{F}_{n-3} branched over a smooth curve D_{n-2} in the linear system $|6C_0 + (4n-8)f|$. R. Fintushel and R. Stern showed that Horikawa surface can be decomposed into

$$H(n) = B_{n-2} \cup D_{n-2} \cup B_{n-2},$$

where B_{n-2} is the complement of a neighborhood of the pair of 2-spheres $(C_0 + (n-2)f)$ and C_0 in \mathbb{F}_{n-3} (Lemma 2.1 in [3]), and they proved that an elliptic surface $E(n)$ is obtained from $H(n)$ by replacing two rational balls B_{n-2} with two configurations C_{n-2} (Lemma 7.3 in [3]). In other words, R. Fintushel and R. Stern proved that Horikawa surface $H(n)$ can be obtained from an elliptic surface $E(n)$ by rationally blowing-down two disjoint configurations C_{n-2} lying in $E(n)$ in smooth category. The aim of this article is to prove that the rational blow-down surgery above can be performed in complex category, which is following.

Proof of Theorem 1.1. Note that \tilde{X}_n is a double covering of \mathbb{F}_n branched over D , and the minimal resolution of two rational double points of type A_1 and A_{2n-9} in \tilde{X}_n is $E(n)$, which is also a double cover of Z_n branched over the proper transform of D . Since the proper transform of D does not meet the contracted linear chain of \mathbb{P}^1 's, we have a double cover of Y_n branched over the image of the proper transform of D by the map ψ . We denote this surface by X_n . Then X_n is the singular surface obtained by contracting two disjoint configurations C_{n-2} from an elliptic surface $E(n)$ and it has two quotient singularities of class T , both are of type $\frac{1}{(n-2)^2}(1, n-3)$. By the fact that the proper transform of D is disjoint from the contracted linear chain of \mathbb{P}^1 's and Lemma 3.4, the map from X_n to Y_n is flat and $H^1(Y_n, \mathcal{O}_{Y_n}(\bar{D}_{Y_n})) = 0$, where \bar{D}_{Y_n} is the image of Δ in Y_n under the contraction C_{n-2} of the linear chain of \mathbb{P}^1 's. Therefore we have the following commutative diagram of maps

$$\begin{array}{ccccc} \tilde{X}_n & \leftarrow & E(n) & \rightarrow & X_n \\ \downarrow & & \downarrow & & \downarrow \\ \mathbb{F}_n & \xleftarrow{\pi} & Z_n & \xrightarrow{\psi} & Y_n \end{array}$$

where all vertical maps are double coverings. Then, by Theorem 2.2 and Proposition 3.2 above, the singular surface X_n has a \mathbb{Q} -Gorenstein smoothing of two quotient singularities all together.

Finally, by applying the standard arguments about Milnor fibers (§5 in [11] or §1 in [13]), we know that a general fiber of a \mathbb{Q} -Gorenstein smoothing of X_n is diffeomorphic to the 4-manifold obtained by rational blow-down of $E(n)$. And we also know that $H(n)$ has one deformation class ([2], Chapter VII). Therefore a general fiber of a \mathbb{Q} -Gorenstein smoothing of X_n is a Horikawa surface $H(n)$ in complex category. \square

REFERENCES

- [1] M. Artin, *Some numerical criteria for contractability of curves on algebraic surfaces*, Amer. J. Math. **84** (1962), 485–496.
- [2] W. Barth, K. Hulek, C. Peters, A. Van de Ven, *Compact complex surfaces*, 2nd ed. Springer-Verlag, Berlin, 2004.
- [3] R. Fintushel and R. Stern, *Rational blowdowns of smooth 4-manifolds*, Jour. Diff. Geom. **46** (1997), 181–235.
- [4] R. Gompf, *A new construction of symplectic manifolds*, Ann. of Math. (2) **142** (1995), 527–595.
- [5] P. Hacking, *Compact moduli of plane curves*, Duke Math. J. **124** (2004), 213–257.
- [6] R. Hartshorne, *Algebraic Geometry*, Springer, 1977.
- [7] B. Hassett, *Stable log surfaces and limits of quartic plane curves*, Manuscripta Math. **100** (1999), 469–497.
- [8] J. Kollár and N.I. Shepherd-Barron, *Threefolds and deformations of surface singularities*, Invent. Math. **91** (1988), 299–338.
- [9] Y. Lee and J. Park, *A simply connected surface of general type with $p_g = 0$ and $K^2 = 2$* , Invent. Math. **170** (2007), 483–505.
- [10] S. Lichtenbaum and M. Schlessinger, *The cotangent complex of a morphism*, Trans. Amer. Math. Soc. **128** (1967), 41–70.
- [11] E. Looijenga and J. Wahl, *Quadratic functions and smoothing surface singularities*, Topology **25** (1986), 261–291.
- [12] M. Manetti, *Normal degenerations of the complex projective plane*, J. Reine Angew. Math. **419** (1991), 89–118.
- [13] M. Manetti, *On the moduli space of diffeomorphic algebraic surfaces*, Invent. Math. **143** (2001), 29–76.
- [14] V. P. Palamodov, *Deformations of complex spaces*, Russian Math. Surveys **31:3** (1976), 129–197.
- [15] I. Reider, *Vector bundles of rank 2 and linear systems on algebraic surfaces*, Ann. Math. **127** (1988), 309–316.
- [16] J. Wahl, *Smoothing of normal surface singularities*, Topology **20** (1981), 219–246.
- [17] J. Wahl, *Elliptic deformations of minimally elliptic singularities*, Math. Ann. **253** (1980), 241–262.

DEPARTMENT OF MATHEMATICS, SOGANG UNIVERSITY, SINSU-DONG, MAPO-GU, SEOUL 121-742, KOREA

E-mail address: ynlee@sogang.ac.kr

DEPARTMENT OF MATHEMATICAL SCIENCES, SEOUL NATIONAL UNIVERSITY, SAN 56-1, SILLIM-DONG, GWANAK-GU, SEOUL 151-747, KOREA

E-mail address: jipark@math.snu.ac.kr